

# Birdwatching Trip Report Colombia



## Choco, Eastern Cordillera & Santa Marta areas

*24 December 2015 - 9 January 2016*

Ariel Brunner, Lu Meldolesi & Pierre Commenville

# Birdwatching trip report

*Choco, Eastern Cordillera & Santa Marta areas*

*24 December 2015 - 9 January 2016*

## Summary

This is the report of the birdwatching trip in Colombia which Ariel Brunner, Lu Meldolesi and Pierre Commenville undertook at the end of the year 2015. Ariel and Lu visited Bahia Solano area in the Choco bioregion for 4 days, then we all had two days birding in the Andes Eastern Cordillera (Chicaque private reserve and Chinganza National Park) and Ariel and Pierre had an eleven day trip in Santa Marta Area (cloud forest, temperate, subtropical, foothills, lowland and wetlands at the western tip of Guajira Peninsula). Ariel spend a few hours on his last day in remnants of the Sabana de Bogota' wetlands (around Parque la Florida). We recorded 434 species and had some amazing birding days, with some amounting to almost 100 species. We had 8 endemic birds restricted to the Sierra Nevada de Santa Marta range, which is an average good result for this endemic bird area recognized as the richest in the World.

## Choco Area

The Choco is one of the most biodiverse, and rainiest places on earth. We took a flight through Medellin to the isolated coastal village of Bahia Solano and then by road to the wonderful El Almejal lodge on the edge of El Valle hamlet. The scenery of humid forest crashing down into the Pacific ocean is truly spectacular. We birded along the main road which crosses both good forest and cleared pastures, took a boat trip up the Rio Valle, spent a long day on the trail leading from El Valle to the Ensenada de Utria National Park and another day visiting the N.P. by boat. We enjoyed the service of a wonderful and extremely friendly local guide Balmes Mosquera.

An unexpected highlight has been the boat ride to Utria N.P. which offered us close up views of Yellowfin tuna and Brown boobies hunting small fish; we had also 3 Band rumped Storm petrel, one of which was terminally exhausted to the

point of being picked up by hand from the water. This bird should not be seen so close to shore and it's occurrence may be linked to this year's strong El Niño. Other highlights have been spectacular observations of Great Jacamar and Baudo Oropendula and a surprisingly high density of both Purple throated Fruitcrow and Black tipped Cotinga. It is worth mentioning that despite being in the dry season, we had heavy rain the whole night, every night, and repeated heavy showers also during the days.

## Andes

The idea of this sequence was to sample some of the very rich variety of birds species in the high altitudes that we will not visit afterwards. We also wanted to see extensive paramo habitat, that we do not know well.

Chicaque private reserve boasted very good surprises and we had a lovely day hiking down to the refuge, located at the basis of the rock cliffs, and letting some of us be driven back to the main entrance at the afternoon. This day-trip from Bogota was a very good option to discover temperate and subtropical humid forests of the Andes. We enjoyed 4 mixed flocks with one of them hosting more than 20 species, encompassing ground to canopy birds. Highlights were good views on Emerald Toucanet, Black-throated Fruit-Eater and Chestnut-crowned Antpitta. Several tanagers species and several brush finches spiced the day, which culminated with a long visit to the hummingbirds feeders at the main entrance, good for the Golden-bellied Starfrontlet and Glowing Pufflegs. Ariel got an impressive Brown-billed Scythebill close to the refuge.



We then visited Chinganza National Park on a day-trip from Bogota. This is worth the journey because this park of high paramo vegetation offers very vast surface of this specific vegetation type of the high Andes. *Espeletia* flowers abound there and the habitat is in very good condition. Weather was not ideal for birding with strong sun and winds (which made however for some stunning mountain scenery) but we managed to get good views of the female of the Green-bearded Helmetcrest and several Flowerpiercer species. Highlight was a Chestnut-breasted Buzzard-Eagle that soared above us for some time. White-tailed Deers are numerous and easy to see. The landscape is fabulous and offers nice views on the Central Cordillera volcanoes.

## Santa Marta Area

We started with three days in Minca, one of the Meccas of birdwatching in Colombia. We birded around Minca and on the road to Pozo Azul, staying at Finca Faunal, a stone throw from this natural pool. Despite the vacation period that brought many Colombians to the freshness of these pools, we had some quiet trails, especially in the early morning. Finca Faunal holds some good habitat despite its limited size (6 hectares only). We had a whole bunch of tanagers, woodpeckers and other colorful species at the feeders and in the garden. On the road to Pozo Azul we saw very well Rufous-breasted and Rufous-and-white Wrens. Hotel Minca has also nice hummingbird feeders crowded with White-necked Jacobins.

We then stayed four nights in Palo Alto, at the entrance of the El Dorado Reserve



managed by ProAves. This lodge that offers basic accommodation has a wonderful garden, very carefully tended by Luis whose parents hold the nearby Tienda de Las Rosas, where we took our meals. This garden is teeming with hummingbirds and we had excellent sightings of Santa Marta Woodstar (males are tricky to see in this season, be patient!) and Blossomcrown there.

Birding long the road at different altitudes, including until the famous San Lorenzo Ridge, was very good to see the specialties. A morning flight of Military Macaws was a special treat. We had



good views of Santa Marta Tapaculo and Santa Marta Foliage Gleaner. In all these days we had more than 50 species per day, denoting an exceptional richness in species of those habitats. Birding on San Lorenzo Ridge in the early morning with views over the snow peaks was noticeably delightful. We had Santa Marta Mountain Tanagers in good numbers and nice Black-throated Tody-Tyrant. All the habitats are still in good conditions and birding along the road makes no problem since the traffic is really low. Night walks were finally great deals for the Band-tailed Guan (the last and the first bird of 2015-2016 passage for us). We also had Kinkajous and Night Monkeys during only short searches.

For wetlands, we headed for Los Camarones, where we stayed for two days at Boca de Camarones. One should take a boat to pay a visit to the stationary group of flamingos, whose number was great when we were there (an approx. of 2,000 individuals). Many local guides offer their service and will also guide you in heronries with great care. We had almost all the possible *Ardeidae* there. Exploring the beaches is good for waders, even it was quite weak for the *Calidris* that should stay in quieter and more remote places during the day. The forest around is also very good and we benefitted from the help of Jose Luis, a local

guide with outstanding knowledge of the birds of these tracts of Tropical Dry Forest, which host god numbers of birds of the Guajira Peninsula.

Unfortunately the forest is being heavily degraded by firewood collection and grazing. We enjoyed the rare sighting (for this region) of Whistling Heron. We also birded song the road to Cari-Cari, a nearby village. Here, we also discovered a mixed-sex group of Dickissel (one male 4 females), a scarce boreal migrant.



Finally we had three days around Tayrona National Park, staying in different places. We explored the rio Don Diego area (along the Tayronaka property), that was good for herons, stream specialists and some good forest birds. We had good sightings of Sooty-capped Hermit, feeding on a mimosa tree. We confirm this observation although the latest edition of ProAves book says this species is unrecorded in Santa Marta areas. Along the river we had good views of several bands of Hawler monkeys. We also visit the Playa de Los Naranjos surroundings that were good for White-bellied Antbird and Lance-tailed Manakin Including displaying mails). Finally we stayed for two nights at Aguas Limpias, to avoid the crowd that press at the National Park gates during the winter holidays. We had a good trail exploring the summit of Lowland forest and moving up to the Foothills forest. We had White-bearded Manakins and several good raptors, including a stunning view of the Short-tailed Hawk. We also had great views of the peculiar Cotton top tamarins.



## La Florida

The last day Ariel used the morning before the flight home to visit Parque La Florida on the outskirts of Bogota. The area holds some tiny and heavily degraded remnants of the wetlands of the original Sabana de Bogota. While the site offers easy views of a number of interesting (including endemic and endangered) species, it was a rather sad close to the trip. One can only hope that some land can be still saved from the encroaching development and properly restored.

## Logistics

We stayed in Hotel Santa Fe Real in Bogota (av Esperanza y 42), that offered good service but is rather badly located, far from the city touristic attractions and nice «barrios». However it was of easy access from the airport and convenient to exit the city.

For the day-trips from Bogota, we hired the service of a private chauffeur for this section, that was really handy and helped to maximise time for birding. We chose Jovani Flores who did a great and discreet service.

For visiting Santa Marta Area, we were for 11 days with Sebastian Ballesteros, who is a young and talented photographer and passionate nature lover. He owns Finca Faunal in Minca from where he has been operating for 5 years. He does an excellent job at the finca, which demonstrates his ability of acting and not only saying. He was of very nice company and everyday did his best to content

us on the best views possible. He knows the surroundings of Santa Marta very well, having spent many days exploring the area and it enables him to be flexible taking into account the season, human pressure and accommodation availability. All in one, Sebastian makes an ideal choice for visiting this region, which we warmly recommend.

For more details do not hesitate to join Ariel [brunner.ariel@birdlife.com](mailto:brunner.ariel@birdlife.com) or Pierre [pierre.commenville@gmail.com](mailto:pierre.commenville@gmail.com)

Mammals and bird lists follow.

Done in Paris & Brussels February 2016



## Mammal list

Red-tailed Squirrel
Venezuelan Red-howler Monkey
Crab-eating Fox
Cottontop Tamarin
Geoffrey's Tamarin
Kinkajou
Night-Monkey
White-tailed Deer
Long tailed weasel



## Bird list

<b>Species</b>	<b>Number of days with sightings</b>
<b>Andean Teal</b>	<b>1</b>
<b>Blue winged teal</b>	<b>1</b>
<b>Masked duck</b>	<b>1</b>
<b>Ruddy duck</b>	<b>1</b>
<b>Black-bellied Whistling-Duck</b>	<b>1</b>
<b>Sickle-winged Guan</b>	<b>2</b>
<b>Andean Guan</b>	<b>1</b>
<b>Band-tailed Guan</b>	<b>4</b>
<b>Grey-headed Chachalaca</b>	<b>1</b>
<b>Rufous-vented Chachalaca</b>	<b>1</b>
<b>Crested Bobwhite</b>	<b>1</b>
<b>Black-fronted Wood-Quail</b>	<b>1</b>
<b>American Flamingo</b>	<b>1</b>
<b>Band-rumped Storm-Petrel</b>	<b>1</b>
<b>Neotropical Cormorant</b>	<b>6</b>
<b>Brown Pelican</b>	<b>5</b>
<b>Brown Booby</b>	<b>1</b>
<b>Magnificent Frigatebird</b>	<b>5</b>
<b>Rufescent Tiger-Heron</b>	<b>1</b>
<b>Black-crowned Night-Heron</b>	<b>2</b>
<b>Yellow-crowned Night-Heron</b>	<b>2</b>
<b>Green Heron</b>	<b>3</b>
<b>Striated Heron</b>	<b>3</b>
<b>Cattle Egret</b>	<b>5</b>
<b>Whistling Heron</b>	<b>2</b>
<b>Great Egret</b>	<b>7</b>
<b>Great Blue Heron</b>	<b>4</b>
<b>Cocoi Heron</b>	<b>3</b>
<b>Tricoloured Heron</b>	<b>2</b>
<b>Reddish Heron</b>	<b>1</b>
<b>Little Blue Heron</b>	<b>5</b>
<b>Snowy Egret</b>	<b>6</b>

<b>Species</b>	<b>Number of days with sightings</b>
<b>White Ibis</b>	<b>2</b>
<b>Scarlet Ibis</b>	<b>1</b>
<b>Bared faced Ibis</b>	<b>1</b>
<b>Roseate Spoonbill</b>	<b>2</b>
<b>Wood Stork</b>	<b>2</b>
<b>Turkey Vulture</b>	<b>15</b>
<b>Lesser Yellow-headed Vulture</b>	<b>1</b>
<b>King Vulture</b>	<b>3</b>
<b>Black Vulture</b>	<b>11</b>
<b>Osprey</b>	<b>4</b>
<b>White tailed Kite</b>	<b>1</b>
<b>Grey-headed Kite</b>	<b>2</b>
<b>Swallow-tailed Kite</b>	<b>1</b>
<b>Savanna Hawk</b>	<b>1</b>
<b>Black-chested Buzzard-Eagle</b>	<b>0</b>
<b>Tiny Hawk</b>	<b>1</b>
<b>Sharp-shinned Hawk</b>	<b>1</b>
<b>Common Black-Hawk</b>	<b>3</b>
<b>White-rumped Hawk</b>	<b>1</b>
<b>Roadside Hawk</b>	<b>4</b>
<b>White Hawk</b>	<b>1</b>
<b>Harris' Hawk</b>	<b>2</b>
<b>White-tailed Hawk</b>	<b>1</b>
<b>Variable Hawk</b>	<b>1</b>
<b>Swainson's Hawk</b>	<b>1</b>
<b>Broad-winged Hawk</b>	<b>5</b>
<b>Zone-tailed Hawk</b>	<b>1</b>
<b>Grey-lined Hawk</b>	<b>4</b>
<b>Short-tailed Hawk</b>	<b>2</b>
<b>Collared Forest Falcon</b>	<b>1</b>
<b>Crested Caracara</b>	<b>4</b>
<b>Yellow-headed Caracara</b>	<b>5</b>

<b>Species</b>	<b>Number of days with sightings</b>
<b>Merlin</b>	<b>1</b>
<b>American Kestrel</b>	<b>3</b>
<b>Bat Falcon</b>	<b>3</b>
<b>Peregrine Falcon</b>	<b>1</b>
<b>Common Gallinule</b>	<b>1</b>
<b>Spot flanked Gallinule</b>	<b>1</b>
<b>American Coot</b>	<b>1</b>
<b>Purple Gallinule</b>	<b>1</b>
<b>Limpkin</b>	<b>1</b>
<b>Grey Plover</b>	<b>3</b>
<b>Southern Lapwing</b>	<b>4</b>
<b>Semipalmated Plover</b>	<b>3</b>
<b>Collared Plover</b>	<b>1</b>
<b>American Oystercatcher</b>	<b>2</b>
<b>Black-necked Stilt</b>	<b>3</b>
<b>Whimbrel</b>	<b>5</b>
<b>Marbled Godwit</b>	<b>1</b>
<b>Ruddy Turnstone</b>	<b>1</b>
<b>Sanderling</b>	<b>2</b>
<b>Western Sandpiper</b>	<b>2</b>
<b>Short-billed Dowitcher</b>	<b>1</b>
<b>Wilson's Snipe</b>	<b>1</b>
<b>Spotted Sandpiper</b>	<b>3</b>
<b>Greater Yellowlegs</b>	<b>3</b>
<b>Lesser Yellowlegs</b>	<b>4</b>
<b>Willet</b>	<b>4</b>
<b>Solitary Sandpiper</b>	<b>4</b>
<b>Wattled Jacana</b>	<b>3</b>
<b>Laughing Gull</b>	<b>2</b>
<b>Gull-billed Tern</b>	<b>1</b>
<b>Sandwich Tern</b>	<b>3</b>
<b>Yellow-billed Tern</b>	<b>1</b>

<b>Species</b>	<b>Number of days with sightings</b>
<b>Elegant Tern</b>	<b>1</b>
<b>Royal Tern</b>	<b>4</b>
<b>Black Skimmer</b>	<b>1</b>
<b>Rock Pigeon</b>	<b>4</b>
<b>Common Ground-Dove</b>	<b>3</b>
<b>Ruddy Ground-Dove</b>	<b>6</b>
<b>Blue Ground-Dove</b>	<b>1</b>
<b>Scaled Dove</b>	<b>2</b>
<b>Band-tailed Pigeon</b>	<b>5</b>
<b>Dusky Pigeon</b>	<b>1</b>
<b>Pale-vented Pigeon</b>	<b>3</b>
<b>Bare-Eyed Pigeon</b>	<b>0</b>
<b>Scaled Pigeon</b>	<b>2</b>
<b>Eared Dove</b>	<b>4</b>
<b>White-tipped Dove</b>	<b>7</b>
<b>Lined Quail-Dove</b>	<b>2</b>
<b>Military Macaw</b>	<b>2</b>
<b>Scarlet-fronted Parakeet</b>	<b>1</b>
<b>Orange-chinned Parakeet</b>	<b>4</b>
<b>Brown-throated Parakeet</b>	<b>2</b>
<b>Green-rumped Parrotlet</b>	<b>2</b>
<b>Blue-headed Parrot</b>	<b>4</b>
<b>Red-lored Parrot</b>	<b>1</b>
<b>Red-billed Parrot</b>	<b>3</b>
<b>Squirrel Cuckoo</b>	<b>4</b>
<b>Ferruginous Pygmy Owl</b>	<b>1</b>
<b>Greater Ani</b>	<b>2</b>
<b>Smooth-billed Ani</b>	<b>3</b>
<b>Groove-billed Ani</b>	<b>2</b>
<b>Common Potoo</b>	<b>1</b>
<b>Lesser Nighthawk</b>	<b>2</b>
<b>Common Pauraque</b>	<b>2</b>

<b>Species</b>	<b>Number of days with sightings</b>
<b>White-tipped Swift</b>	<b>1</b>
<b>White-collared Swift</b>	<b>4</b>
<b>White-necked Jacobin</b>	<b>2</b>
<b>Rufous-breasted Hermit</b>	<b>2</b>
<b>White-tipped Sicklebill</b>	<b>1</b>
<b>Pale-bellied Hermit</b>	<b>1</b>
<b>Sooty-capped Hermit</b>	<b>1</b>
<b>Long-billed Hermit</b>	<b>1</b>
<b>Sparkling Violetear</b>	<b>4</b>
<b>Green Violetear</b>	<b>6</b>
<b>Purple-crowned Fairy</b>	<b>1</b>
<b>Ruby-topaz Hummingbird</b>	<b>1</b>
<b>Black-Throated Mango</b>	<b>2</b>
<b>Green-bearded Helmetcrest</b>	<b>1</b>
<b>Tourmaline Sunangel</b>	<b>1</b>
<b>Tyrian Metaltail</b>	<b>3</b>
<b>Glowing Puffleg</b>	<b>1</b>
<b>Blossomcrown</b>	<b>2</b>
<b>Gorgeted Woodstar</b>	<b>1</b>
<b>Santa Marta Woodstar</b>	<b>4</b>
<b>Red-billed Emerald</b>	<b>3</b>
<b>Coppery Emerald</b>	<b>1</b>
<b>Collared Inca</b>	<b>1</b>
<b>Golden-bellied Starfrontlet</b>	<b>1</b>
<b>Buff-tailed Coronet</b>	<b>1</b>
<b>White-tailed Starfrontlet</b>	<b>1</b>
<b>Long-billed Starthroat</b>	<b>1</b>
<b>White-vented Plumeleteer</b>	<b>2</b>
<b>Rufous-tailed Hummingbird</b>	<b>6</b>
<b>Blue-chested Hummingbird</b>	<b>1</b>
<b>White-chinned Sapphire</b>	<b>1</b>
<b>Steely-vented Hummingbird</b>	<b>3</b>



<b>Species</b>	<b>Number of days with sightings</b>
<b>Buffy Hummingbird</b>	<b>1</b>
<b>Crowned Woodnymph</b>	<b>5</b>
<b>Shining-green Hummingbird</b>	<b>2</b>
<b>White-tailed Trogon</b>	<b>1</b>
<b>Gartered Trogon</b>	<b>3</b>
<b>Masked Trogon</b>	<b>4</b>
<b>Ringed Kingfisher</b>	<b>3</b>
<b>Belted Kingfisher</b>	<b>2</b>
<b>Amazon Kingfisher</b>	<b>2</b>
<b>Green Kingfisher</b>	<b>2</b>
<b>Green-and-rufous Kingfisher</b>	<b>1</b>
<b>Whooping Motmot</b>	<b>3</b>
<b>Rufous-tailed Jacamar</b>	<b>6</b>
<b>Great Jacamar</b>	<b>1</b>
<b>Black-breasted Puffbird</b>	<b>1</b>
<b>Pied Puffbird</b>	<b>1</b>
<b>Russet-throated Puffbird</b>	<b>4</b>
<b>White-whiskered Puffbird</b>	<b>1</b>
<b>Spot-crowned Barbet</b>	<b>1</b>
<b>Five-coloured Barbet</b>	<b>1</b>
<b>Chestnut-mandibled Toucan</b>	<b>1</b>
<b>Choco Toucan</b>	<b>1</b>
<b>Keel-billed Toucan</b>	<b>7</b>
<b>Emerald Toucanet</b>	<b>3</b>
<b>Collared Aracari</b>	<b>5</b>
<b>Scaled Piculet</b>	<b>2</b>
<b>Chestnut Piculet</b>	<b>1</b>
<b>Black-cheeked Woodpecker</b>	<b>1</b>
<b>Smoky-brown Woodpecker</b>	<b>2</b>
<b>Red-crowned Woodpecker</b>	<b>6</b>
<b>Crimson-mantled Woodpecker</b>	<b>1</b>
<b>Golden-olive Woodpecker</b>	<b>4</b>

<b>Species</b>	<b>Number of days with sightings</b>
<b>Spot-breasted Woodpecker</b>	<b>1</b>
<b>Cinnamon Woodpecker</b>	<b>1</b>
<b>Lineated Woodpecker</b>	<b>2</b>
<b>Crimson-crested Woodpecker</b>	<b>2</b>
<b>Barred Antshrike</b>	<b>3</b>
<b>Black-crested Antshrike</b>	<b>2</b>
<b>Black-crowned Antshrike</b>	<b>1</b>
<b>Santa Marta Antbird</b>	<b>1</b>
<b>Checker-throated Antwren</b>	<b>1</b>
<b>Moustached Antwren</b>	<b>1</b>
<b>Pacific Antwren</b>	<b>1</b>
<b>White-flanked Antwren</b>	<b>1</b>
<b>Slaty Antwren</b>	<b>1</b>
<b>White-fringed Antwren</b>	<b>2</b>
<b>Dusky Antbird</b>	<b>1</b>
<b>White-bellied Antbird</b>	<b>2</b>
<b>Chestnut-backed Antbird</b>	<b>1</b>
<b>Spotted Antbird</b>	<b>1</b>
<b>Rufous Antpitta</b>	<b>1</b>
<b>Chestnut-crowned Antpitta</b>	<b>1</b>
<b>Rufous-breasted Antpitta</b>	<b>1</b>
<b>Black-headed Anttrush</b>	<b>1</b>
<b>Santa Marta Tapaculo</b>	<b>1</b>
<b>Blackish Tapaculo</b>	<b>1</b>
<b>Grey-throated Leaftosser</b>	<b>2</b>
<b>Brown-billed Scythebill</b>	<b>1</b>
<b>Montane Woodcreeper</b>	<b>5</b>
<b>Black-banded Woodcreeper</b>	<b>2</b>
<b>Strong-billed Woodcreeper</b>	<b>2</b>
<b>Cocoa Woodcreeper</b>	<b>3</b>
<b>Streak-headed Woodcreeper</b>	<b>3</b>
<b>Black-striped Woodcreeper</b>	<b>1</b>

<b>Species</b>	<b>Number of days with sightings</b>
<b>Spotted Woodcreeper</b>	<b>1</b>
<b>Straight-billed Woodcreeper</b>	<b>5</b>
<b>Plain Xenops</b>	<b>2</b>
<b>Streaked Xenops</b>	<b>2</b>
<b>Montane Foliage-gleaner</b>	<b>5</b>
<b>Santa Marta Foliage-gleaner</b>	<b>1</b>
<b>Rusty-winged Barbtail</b>	<b>1</b>
<b>Spotted Barbtail</b>	<b>1</b>
<b>Rufous Spinetail</b>	<b>1</b>
<b>Streak-capped Spinetail</b>	<b>1</b>
<b>Pale-breasted Spinetail</b>	<b>1</b>
<b>Rusty-headed Spinetail</b>	<b>1</b>
<b>White-whiskered Spinetail</b>	<b>3</b>
<b>Pale-legged Hornero</b>	<b>1</b>
<b>Sooty-headed Tyrannulet</b>	<b>1</b>
<b>Yellow-bellied Elaenia</b>	<b>5</b>
<b>Lesser Elaenia</b>	<b>1</b>
<b>Brown-capped Tyrannulet</b>	<b>1</b>
<b>White-throated Tyrannulet</b>	<b>1</b>
<b>Sepia-capped Flycatcher</b>	<b>1</b>
<b>Forest Elaenia</b>	<b>2</b>
<b>Coopman's Tyrannulet</b>	<b>3</b>
<b>Torrent Tyrannulet</b>	<b>1</b>
<b>Black-capped Tyrannulet</b>	<b>2</b>
<b>Olive-striped Fruit-tyrant</b>	<b>7</b>
<b>Streak-necked Fruit-tyrant</b>	<b>1</b>
<b>Ochre-bellied Fruit-tyrant</b>	<b>2</b>
<b>Northern Scrub-Flycatcher</b>	<b>1</b>
<b>Slender-billed Inezia</b>	<b>2</b>
<b>Southern Bentbill</b>	<b>1</b>
<b>Black-capped Pygmy-Tyrant</b>	<b>1</b>
<b>Pale-eyed Pygmy-Tyrant</b>	<b>1</b>

<b>Species</b>	<b>Number of days with sightings</b>
<b>Pearly-vented Tody-Tyrant</b>	<b>1</b>
<b>Black-throated Tody-Tyrant</b>	<b>1</b>
<b>Slate-headed Tody-Flycatcher</b>	<b>1</b>
<b>Yellow-margined Flycatcher</b>	<b>1</b>
<b>Yellow-breasted Flycatcher</b>	<b>1</b>
<b>Cinnamon Flycatcher</b>	<b>6</b>
<b>Handsome Flycatcher</b>	<b>1</b>
<b>Vermilion Flycatcher</b>	<b>2</b>
<b>Plain-capped Ground-Tyrant</b>	<b>1</b>
<b>Pied Water-Tyrant</b>	<b>1</b>
<b>Slaty-backed Chat-Tyrant</b>	<b>1</b>
<b>Brown-backed Chat-Tyrant</b>	<b>1</b>
<b>Long-tailed Tyrant</b>	<b>1</b>
<b>Cattle Tyrant</b>	<b>2</b>
<b>Rusty-margined Flycatcher</b>	<b>1</b>
<b>Tropical Pewee</b>	<b>2</b>
<b>Olive-sided Flycatcher</b>	<b>3</b>
<b>Social Flycatcher</b>	<b>4</b>
<b>Boat-billed Flycatcher</b>	<b>3</b>
<b>Great Kiskadee</b>	<b>3</b>
<b>Golden-crowned Flycatcher</b>	<b>1</b>
<b>Streaked Flycatcher</b>	<b>5</b>
<b>Piratic Flycatcher</b>	<b>1</b>
<b>Choco Syristes</b>	<b>1</b>
<b>Grey Kingbird</b>	<b>4</b>
<b>Tropical Kingbird</b>	<b>9</b>
<b>Fork-tailed Flycatcher</b>	<b>1</b>
<b>Brown-capped Flycatcher</b>	<b>2</b>
<b>Pale-edged Flycatcher</b>	<b>1</b>
<b>Dusky-capped Flycatcher</b>	<b>3</b>
<b>Panama Flycatcher</b>	<b>3</b>
<b>Green-and-black Fruiteater</b>	<b>1</b>

<b>Species</b>	<b>Number of days with sightings</b>
<b>Golden-breasted Fruiteater</b>	<b>2</b>
<b>Purple-throated Fruit-Crow</b>	<b>1</b>
<b>Back-tipped Cotinga</b>	<b>1</b>
<b>White-bearded Manakin</b>	<b>3</b>
<b>Lance-tailed Manakin</b>	<b>2</b>
<b>Golden-collared Manakin</b>	<b>1</b>
<b>Blue-crowned Manakin</b>	<b>1</b>
<b>Masked Tityra</b>	<b>4</b>
<b>Black-crowned Tityra</b>	<b>1</b>
<b>Cinnamon Becard</b>	<b>2</b>
<b>Yellow-throated Vireo</b>	<b>1</b>
<b>Brown-capped Vireo</b>	<b>7</b>
<b>Yellow-green Vireo</b>	<b>1</b>
<b>Golden-fronted Greenlet</b>	<b>2</b>
<b>Scrub Greenlet</b>	<b>1</b>
<b>Black-chested Jay</b>	<b>7</b>
<b>Southern Rough-winged Swallow</b>	<b>5</b>
<b>Brown-chested Martin</b>	<b>1</b>
<b>Grey-breasted Martin</b>	<b>5</b>
<b>Sand Martin</b>	<b>1</b>
<b>Barn Swallow</b>	<b>1</b>
<b>Scaly-breasted Wren</b>	<b>1</b>
<b>House Wren</b>	<b>7</b>
<b>Apolinar's Wren</b>	<b>1</b>
<b>White-headed Wren</b>	<b>1</b>
<b>Bicoloured Wren</b>	<b>6</b>
<b>Rufous-breasted Wren</b>	<b>1</b>
<b>Rufous-and-white Wren</b>	<b>2</b>
<b>Buff-breasted Wren</b>	<b>3</b>
<b>Gray-breasted Wood Wren</b>	<b>1</b>
<b>Santa Marta Wood Wren</b>	<b>3</b>

<b>Species</b>	<b>Number of days with sightings</b>
<b>Bang's Wood Wren</b>	<b>2</b>
<b>Long-billed Gnatwren</b>	<b>2</b>
<b>Tropical Gnatcatcher</b>	<b>3</b>
<b>Orange-billed Nightingale Thrush</b>	<b>1</b>
<b>Grey-cheeked Thrush</b>	<b>1</b>
<b>Swainson's Thrush</b>	<b>3</b>
<b>Great Thrush</b>	<b>2</b>
<b>Yellow-legged Thrush</b>	<b>4</b>
<b>Pale-breasted Thrush</b>	<b>5</b>
<b>Clay-coloured Thrush</b>	<b>3</b>
<b>Black-hooded Thrush</b>	<b>4</b>
<b>White-necked Thrush</b>	<b>3</b>
<b>Tropical Mockingbird</b>	<b>4</b>
<b>Oleaginous Hemispingus</b>	<b>1</b>
<b>Black-eared Hemispingus</b>	<b>1</b>
<b>Grass-green Tanager</b>	<b>1</b>
<b>Grey-headed Tanager</b>	<b>1</b>
<b>Scarlet-bellied Mountain Tanager</b>	<b>1</b>
<b>Santa Marta Mountain Tanager</b>	<b>2</b>
<b>Tawny-crested Tanager</b>	<b>1</b>
<b>White-lined Tanager</b>	<b>2</b>
<b>Grey-headed Tanager</b>	<b>2</b>
<b>Crimson-backed Tanager</b>	<b>7</b>
<b>Lemon-rumped Tanager</b>	<b>1</b>
<b>Palm Tanager</b>	<b>6</b>
<b>Blue-grey Tanager</b>	<b>8</b>
<b>Glaucous Tanager</b>	<b>1</b>
<b>Blue-capped Tanager</b>	<b>3</b>
<b>Beryl-spangled Tanager</b>	<b>1</b>
<b>Plain-coloured Tanager</b>	<b>1</b>
<b>Flame-faced Tanager</b>	<b>1</b>

<b>Species</b>	<b>Number of days with sightings</b>
<b>Black-headed Tanager</b>	<b>4</b>
<b>Black-capped Tanager</b>	<b>5</b>
<b>Bay-headed Tanager</b>	<b>8</b>
<b>Swallow Tanager</b>	<b>3</b>
<b>Scarlet-tighed Dacnis</b>	<b>1</b>
<b>Blue Dacnis</b>	<b>3</b>
<b>Purple Honeycreeper</b>	<b>2</b>
<b>Red-legged Honeycreeper</b>	<b>3</b>
<b>Glossy Flowerpiercer</b>	<b>1</b>
<b>Rusty Flowerpiercer</b>	<b>3</b>
<b>White-sided Flowerpiercer</b>	<b>6</b>
<b>Bluish Flowerpiercer</b>	<b>1</b>
<b>Masked Flowerpiercer</b>	<b>1</b>
<b>Plushcap</b>	<b>1</b>
<b>Variable Seedeater</b>	<b>1</b>
<b>Yellow-bellied Seedeater</b>	<b>3</b>
<b>Grey Seedeater</b>	<b>1</b>
<b>Pileated Finch</b>	<b>2</b>
<b>Bananaquit</b>	<b>4</b>
<b>Black-faced Grassquit</b>	<b>2</b>
<b>Blue-black Grassquit</b>	<b>1</b>
<b>Dusky-faced Tanager</b>	<b>1</b>
<b>Buff-throated Saltator</b>	<b>6</b>
<b>Greyish Saltator</b>	<b>6</b>
<b>Streaked Saltator</b>	<b>1</b>
<b>Orange-billed Sparrow</b>	<b>1</b>
<b>Golden-winged Sparrow</b>	<b>1</b>
<b>Rufous-collared Sparrow</b>	<b>6</b>
<b>Yellow-throated Chlorospingus</b>	<b>1</b>
<b>Common Chlorospingus</b>	<b>1</b>
<b>Grey-browed Brushfinch</b>	<b>1</b>
<b>Moustached Bushfinch</b>	<b>1</b>

<b>Species</b>	<b>Number of days with sightings</b>
<b>Chestnut-capped Brushfinch</b>	<b>1</b>
<b>Colombian Brushfinch</b>	<b>1</b>
<b>Santa Marta Brushfinch</b>	<b>4</b>
<b>Summer Tanager</b>	<b>6</b>
<b>Lemon-spectacled Tanager</b>	<b>1</b>
<b>Vermilion Cardinal</b>	<b>3</b>
<b>Golden-bellied Grosbeak</b>	<b>3</b>
<b>Rose-breasted Grosbeak</b>	<b>2</b>
<b>Blue-black Grosbeak</b>	<b>3</b>
<b>Dickissel</b>	<b>1</b>
<b>Northern Waterthrush</b>	<b>3</b>
<b>Black-and-white Warbler</b>	<b>7</b>
<b>Protonothary Warbler</b>	<b>4</b>
<b>Tennessee Warbler</b>	<b>8</b>
<b>Bay-breasted Warbler</b>	<b>1</b>
<b>Blackburnian Warbler</b>	<b>8</b>
<b>Yellow Warbler</b>	<b>3</b>
<b>Chestnut-sided Warbler</b>	<b>1</b>
<b>Santa Marta Warbler</b>	<b>1</b>
<b>Three-striped Warbler</b>	<b>1</b>
<b>American Redstart</b>	<b>5</b>
<b>Slate-throated Whitestart</b>	<b>6</b>
<b>Golden-fronted Redstart</b>	<b>1</b>
<b>Yellow-crowned Whitestart</b>	<b>2</b>
<b>Rufous-capped Warbler</b>	<b>2</b>
<b>Golden-crowned Warbler</b>	<b>2</b>
<b>White-lored Warbler</b>	<b>4</b>
<b>Yellow-rumped Cacique</b>	<b>0</b>
<b>Yellow-billed Cacique</b>	<b>1</b>
<b>Chestnut-headed Oropendola</b>	<b>1</b>
<b>Crested Oropendola</b>	<b>6</b>
<b>Baudo Oropendola</b>	<b>1</b>



<b>Species</b>	<b>Number of days with sightings</b>
<b>Scarlet-rumped Cacique</b>	<b>1</b>
<b>Orange-crowned Oriole</b>	<b>1</b>
<b>Yellow-backed Oriole</b>	<b>1</b>
<b>Yellow Oriole</b>	<b>3</b>
<b>Baltimore Oriole</b>	<b>1</b>
<b>Yellow headed Blackbird</b>	<b>1</b>
<b>Shiny Cowbird</b>	<b>2</b>
<b>Giant Cowbird</b>	<b>1</b>
<b>Carib Grackle</b>	<b>3</b>
<b>Great-tailed Grackle</b>	<b>4</b>
<b>Red-breasted Blackbird</b>	<b>1</b>
<b>Eastern Meadowlark</b>	<b>1</b>
<b>Andean siskin</b>	<b>3</b>
<b>Lesser Goldfinch</b>	<b>4</b>
<b>Trinidad Euphonia</b>	<b>2</b>
<b>Thick-billed Euphonia</b>	<b>5</b>
<b>Fulvous-vented Euphonia</b>	<b>1</b>
<b>White-vented Euphonia</b>	<b>1</b>
<b>Blue-naped Chorophonia</b>	<b>4</b>
<b>House Sparrow</b>	<b>1</b>
<b>TOTAL NB</b>	<b>434</b>